arXiv:1210.1090v1 [cond-mat.mes-hall] 3 Oct 2012

Waves of spin-current in magnetized dielectrics

P. A. Andreev*
Department of General Physics, Faculty of Physics,
Moscow State University, Moscow, Russian Federation.

L. S. Kuz'menkov†
Department of Theoretical Physics, Physics Faculty,
Moscow State University, Moscow, Russian Federation.

(Dated: October 4, 2012)

Spin-current is the important physical quantity in present day spintronics and it might be very usefull in the physics of quantum plasma of spinning particles. Thus it is important to have an equation of spin-current dynamical evolution. This equation naturally appears as a part of a set of the quantum hydrodynamics (QHD) equations. Consequently, we present the set of the QHD equations derived from the many-particle microscopic Schrodinger equation, which consists of the continuity equation, the Euler equation, the Bloch equation and equation of spin-current evolution. We use these equations to study dispersion of collective excitations in three dimensional samples of magnetized dielectrics. We show that dynamics of spin-current leads to formation of new type of collective excitations in magnetized dielectrics, which we called spin-current waves.

I. INTRODUCTION

Spin current is very important characteristic, which useful for description such processes as spin injection [1] and other process spin transport involved. All of them accumulated in a grate field of physics-spintronics [2], [3], which has been involved graphene using [4], [5]. It has been expected that silicon spintronics has potential to change information technology, this possibility discussed in Ref. [6]. Processes of spin transport and relaxation in grapheme have also been considered recently [7]. There are different methods of the spic-current generation, such as spin pumping by sound waves [8], optical spin injection [9], and at using of junctions as Polarizers [10], spin flip in the result of electron interaction with electromagnetic wave [11]. Spin-polarized currents are used in spin laser's development [12] and in spin-diode structures [13]. It is interesting to admit that in some devises the spin current exhibits the sine wave-like behavior [14]. New modification of spin field effect transistors have been suggested (see for example [15], [16]), one of the spintronic devices utilize the electron's spin property in addition to its charge property. Effect in junction play key role in spintronic devises. Recently, an interesting effect was observed experimentally, it is the spin-dependent Peltier effect [17], which is based on the ability of the spin-up and spin-down channels to transport heat independently.

Thus, the spin current and it's evolution equation are in the center of attention of this paper. Let's admit that the spin current for many-particle system of charged spinning particles appears in the QHD [18], [19], and in this paper we derive equation for spin current evolution due to QHD method. Recently, an analogous equation was derived for graphene which has also been obtained by the QHD method. This definition and corresponding equations appears in semi-relativistic theory. The spin current in a tensor form have been universally defined according to the quantum electrodynamics in Ref. [20].

The spin current evolution also gives influence on the spin waves, which are well-known phenomenon in many different physical systems, first of all they exist in ferromagnetic and other structures with strong magnetization. For example quantum dots show interesting spin waves behavior [21]- [25]. Dynamics of magnetic moments in plasma has also been studied. In Ref. [26] existence of the self-consistent spin waves in magnetized plasma was showed. Thus new branches of wave dispersion appear in magnetized plasma due to dynamics of magnetic moment of electrons and ions. It was demonstrated by means of the quantum hydrodynamics (QHD) method [27]- [30]. Later, the generalization of the Vlasov's equation for the plasma of spinning charged particles was used to study the same problem [31]. Magnetic moment dynamics also leads to existence of the effect of resonances interaction of the neutron beam with the magnetized plasma, which gives new method of generation of waves in magnetized plasma [19], [32], [33]. Method of the QHD has become very popular and powerfull method of studying influence of magnetic moment dynamics on various processes in magnetized plasma [19], [34]- [38]. Usually the set of the QHD equations for spinning charged particles consists of three equations for evolution of material fields, these are the continuity equation for particles concentration, the Euler equation for the velocity field, and the generalized Bloch equation for magnetic moment evolution, and the Maxwell's equations for the electromagnetic field description.

If we derive the QHD equations from the Pauli equation for one-particle in an external field we find three equation of material field evolution mentioned above. However, if we have deal with many-particle system the set of the QHD equations contains some new functions, for example kinetic pressure $p^{\alpha\beta}$ caused by thermal motion of particles, and spin current $J^{\alpha\beta}$. Two-particle macroscopic functions also appear in terms describing inter-particle interaction. In the self-consistent field approximation, containing no contribution of exchange interaction the terms describing interaction have same form as the terms caused by external fields.

To continue comparison of the many-particle QHD with the one-particle one we admit that in one particle case $p^{\alpha\beta} = 0$ and $J^{\alpha\beta} = M^{\alpha}v^{\beta}$ where M^{α} is the density of magnetic moments, and v^{β} is the velocity field. We can also notice that in one-particle case the kinetic energy field ε easily connected with the particle concentration n, magnetization M^{α} and velocity field v^{β}, so we have $\varepsilon = mnv^2/2$. For many-particle system we have that $p^{\alpha\beta}$, and $J^{\alpha\beta}$ and ε are independent material fields additional to n, v^{α}, and M^{α}. In many-particle system $J^{\alpha\beta}$ and ε partly connect with n, v^{α}, and M^{α}. Let's make an example to describe last statement. For the energy we have $\varepsilon = mnv^2/2 + n\epsilon$, where the last term corresponds to internal energy caused by thermal motion of particles. A set of QHD equations including the energy evolution equation and non-zero thermal pressure in the Euler equation can be derived from a quantum kinetic equation, but derivation of the quantum kinetic equation needs some additional assumption to construct distribution function. During several last decades a lot of different ways of derivation of quantum kinetic equation was suggested, but there are a lot of open questions in this field. Derivation of kinetic equation via the Wigner distribution function is the most popular and actively used in recent publications. However we keep developing the method of many-particle quantum hydrodynamics which is more direct way of derivation of equations for collective quantum dynamics.

QHD description of self-consistent spin waves in a system of neutral particles in one-, two-, and three-dimensional dielectrics was described in Ref. [39].

The method of the QHD can also be used for systems of neutral particles with magnetic moment. Such systems are more preferable to demonstrate results given by the QHD method at description of physical effects caused by evolution of magnetic moments. The QHD method allows us to derive equations of collective dynamics for various physical systems, it has been done for several physical systems, which are

Thus, to make more detailed study of magnetic moment dynamics we going to derive equation evolution for magnetic moment current (or spin current) $J^{\alpha\beta}$ and study it's influence on the spin waves dispersion.

We consider two kind of equilibrium states. One of them corresponds to the case when equilibrium spin current $J^{\alpha\beta}$ equals to zero, in second case we consider non zero equilibrium spin current $J^{\alpha\beta}$, but we suppose that equilibrium velocity field equals to zero. Such structure might be realized by means two currents (flows of neutral particles) directed in opposite directions and having opposite equilibrium spin.

Presented here results are also important for the physics of magnetized ultracold quantum gases. Used where models are equivalent to the first three equations of the QHD, they are the continuity equation, the Euler equation, and the Bloch equation [40], [41].

We also need to accent the fact that the many-particle QHD method has been used for different physical systems, thus the sets of the QHD equations have been made for graphene [42], neutral ultracold quantum gases [43], Bose-Einstein condensate of excitons in graphene [44], along with physics of plasma described above.

This paper is organized as follows. In Sec. II we present and describe the set of the QHD equations derives in the paper. In Sec. III dispersion of spin waves is considered, a contribution of the spin-current in spin wave properties is studied. In Sec. IV brief summary of obtained results are presented.

II. THE MODEL

The many-particle QHD equations are derived from the microscopic many-particle Schrodinger equation. The structure of the QHD equations depends on the explicit form of the Hamiltonian of considered system of particles. We do not described here the method of derivation of the QHD equations, a lot of paper are dedicated to this topic [27]- [30]. However, to be certain we present the Hamiltonian used for derivation of equations presented below.

$$\hat{H} = \sum_{p}\biggl(\frac{1}{2m_p}D_p^2 + e_p\varphi_p^{ext} - \gamma_p\sigma_p^{\alpha}B_{p(ext)}^{\alpha}\biggr)$$

$$+\frac{1}{2}\sum_{p,n\neq p}(e_pe_nG_{pn} - \gamma_p\gamma_nG_{pn}^{\alpha\beta}\sigma_p^{\alpha}\sigma_n^{\beta}), \qquad (1)$$

where $D_n^{\alpha} = -\imath\hbar\partial_n^{\alpha} - e_nA_{n,ext}^{\alpha}/c$, $\varphi_{n,ext}$, $A_{n,ext}^{\alpha}$ - the potentials of the external electromagnetic field, $\partial_n^{\alpha} = \nabla_n^{\alpha}$ is the derivatives on space variables, and $G_{nk} = 1/r_{nk}$, - is the Green functions of the Coulomb interaction, $G_{pn}^{\alpha\beta} = 4\pi\delta^{\alpha\beta}\delta(\mathbf{r}_{pn}) + \partial_p^{\alpha}\partial_p^{\beta}(1/r_{pn})$ is the Green function of spin-spin interaction, γ_p - is the gyromagnetic ratio, σ_p^{α} is the Pauli matrix, a commutation relations for them is

$$[\sigma_p^{\alpha}, \sigma_n^{\beta}] = 2\imath\delta_{pn}\varepsilon^{\alpha\beta\gamma}\sigma_p^{\gamma},$$

e_n, m_n-are the charge and the mass of particle, $\hbar$-is the Planck constant and c is the speed of light. For electrons γ_p reads $\gamma_p = e_p\hbar/(2m_pc)$, $e_p = -|e|$.

Using evident form of the Hamiltonian, we obtain the chain of equations, we break chain of the QHD equations including four equations only. These are equations evolution for concentration n, velocity field v^{β}, density of magnetic moment or spin M^{α}, and spin-current $J^{\alpha\beta}$. First equation of the set of the QHD equation is the continuity equation:

$$\partial_t n + \nabla(n\mathbf{v}) = 0. \qquad (2)$$

Differentiating the function of current with respect to time, we obtain the momentum balance equations, this equation is an analog of the Euler equation:

$$mn(\partial_t + v^\beta\nabla^\beta)v^\alpha + \partial_\beta p^{\alpha\beta}$$

$$-\frac{\hbar^2}{4m}\partial^\alpha\triangle n + \frac{\hbar^2}{4m}\partial^\beta\left(\frac{\partial^\alpha n\cdot\partial^\beta n}{n}\right)$$

$$= enE^\alpha + \frac{e}{c}\varepsilon^{\alpha\beta\gamma}nv^\beta B^\gamma + M^\beta\nabla^\alpha B^\beta, \tag{3}$$

where **E** and **B** are the electric and magnetic fields, **M** is the density of magnetic moments, $\varepsilon^{\alpha\beta\gamma}$ - is the antisymmetric symbol (the Levi-Civita symbol), $p^{\alpha\beta}$ is the kinetic pressure tensor. The momentum balance equation (3) has usual form, we see that evolution of the velocity field caused by momentum current on thermal velocities $p^{\alpha\beta}$, specific for quantum kinematics quantum Bohm potential (two terms in the left-hand side of equation (3), which proportional to $\hbar^2$), and interaction, which is presented in the right-hand side of the momentum balance equation (3). We derive this equation for charged spinning particles, thus the force field contains the density of the Lorentz force describing action of electromagnetic field on charges presented by two first terms and force acting on magnetic moment density from magnetic field presented by the last term.

The equation of evolution of the magnetic moment is:

$$\partial_t M^\alpha + \nabla^\beta J_M^{\alpha\beta} = \frac{2\gamma}{\hbar}\varepsilon^{\alpha\beta\gamma}M^\beta B^\gamma. \tag{4}$$

This equation is an analog of the Bloch equation. From equation (4) we see that evolution of magnetic moment density caused by both the spin current $J^{\alpha\beta}$ and interaction of the magnetic moments with the magnetic field. Charge of particles gives no interference in dynamics of magnetic moment density.

Next equation is the equation of spin-current evolution. The spin-current appears in the Bloch equation, and if we include the spin-orbit and the spin-current interaction we get that the spin-current gives contribution in the force field in the Euler equation (3), for example see Ref. [19]

$$\partial_t J^{\alpha\beta} + \partial^\gamma(J^{\alpha\beta}v^\gamma)$$

$$= \frac{\gamma^2}{m}n\partial^\beta B^\alpha - \frac{2\gamma}{\hbar}\varepsilon^{\alpha\gamma\delta}B^\gamma J^{\delta\beta}. \tag{5}$$

for current of the spin-current $J^{\alpha\beta\gamma}$ which emerges in the second term in the left-hand side of equation (5) we have used approximate formula $J^{\alpha\beta\gamma} = J^{\alpha\beta}v^\gamma$. In general case $J^{\alpha\beta\gamma}$ has more complex structure and contains additional contribution of both the thermal motion and quantum kinematics. The last one shows in the form of a term analogous to the quantum Bohm potential. We do not consider these contributions and pay attention to the spin current evolution caused by inter-particle interaction. We should pay special attention to equation (5) because all this paper is dedicated to consideration of the influence of the spin-current evolution on dynamics of particles system. Equation (5) is presented for the chargeless spinning particles, and this form will be used in the paper. However, now we present it for the charged spinning particles

$$\partial_t J^{\alpha\beta} + \partial^\gamma(J^{\alpha\beta}v^\gamma) = \frac{e}{m}M^\alpha E^\beta$$

$$+\frac{e}{mc}\gamma\varepsilon^{\beta\gamma\delta}J^{\alpha\gamma}B^\delta + \frac{\gamma^2}{m}n\partial^\beta B^\alpha - \frac{2\gamma}{\hbar}\varepsilon^{\alpha\gamma\delta}B^\gamma J^{\delta\beta}. \tag{6}$$

Equations (2)-(5) take place for each species of particles. The electric **E** and magnetic **B** fields appearing in the equations (2)-(5) are caused by charges, electric currents, and magnetic moments of medium and satisfy to the Maxwell's equations. Thus the QHD equations for each species of particles connect by means of the Maxwell's equations:

$$\nabla\mathbf{B} = 0,\ \ \nabla\mathbf{E} = 4\pi\textstyle\sum_a e_a n_a\ ,$$

$$\nabla\times\mathbf{E}(\mathbf{r},t) = 0,$$

$$\nabla\times\mathbf{B}(\mathbf{r},t) = \frac{4\pi}{c}\sum_a e_a n_a\mathbf{v}_a + 4\pi\sum_a\nabla\times\mathbf{M}_a, \tag{7}$$

where subindex "a" describes the species of particles. Presented here Maxwell's equations (7) do not contain time derivatives of the electric and the magnetic fields, because we have derived the QHD equations from non-relativistic theory.

III. DISPERSION EQUATION

We consider collective eigen-waves in a system of neutral spinning particles being in an external uniform magnetic field. So, we have deal with the paramagnetic and diamagnetic dielectrics. There are two fundamental collective excitations in such physical systems, they are sound waves and spin waves. Following to the QHD description of three dimensional magnetized dielectrics [39] we can admit that in such systems there is one type of the spin waves with the constant eigen-frequency $\omega = 2\gamma B_0/\hbar$, where . Here we have no dependency on wave vector, consequently the group velocity $\partial\omega/\partial k$ of these waves equal to zero.

We are interested in interference of the spin-current evolution on dispersion properties of the medium. To find the dispersion dependence of eigen-waves in the described system we consider small amplitude excitations around equilibrium state of the medium. We consider two different equilibrium states. In the first case we suppose that equilibrium spin current equal to zero, and in the second case we consider medium with an equilibrium spin current under condition that equilibrium velocity field is equal to zero.

Here we describe hydrodynamic variables as the sum of an equilibrium part and a small perturbation

$$n_a = n_{0a} + \delta n_a,\ \ \mathbf{E} = 0 + \mathbf{E}\ ,$$

$$\mathbf{B} = B_0\mathbf{e}_z + \delta\mathbf{B},\ \ \mathbf{v}_a = 0 + \mathbf{v}_a\ ,$$

$$M_a^\alpha = M_{0a}^\alpha + \delta M_a^\alpha,\ M_{0a}^\alpha = \chi_a B_0^\alpha,\ J^{\alpha\beta} = J_0^{\alpha\beta} + \delta J^{\alpha\beta},$$

$$n_{0e} = n_{0i},\ p_a^{\alpha\beta} = p_a\delta^{\alpha\beta},\ \delta p_a = m_a v_{sa}^2 \delta n_a, \qquad (8)$$

where $\delta^{\alpha\beta}$- is the Kronecker symbol, $\chi_a = \kappa_a/\nu_a$ is the ratio between the equilibrium magnetic susceptibility κ_a and magnetic permeability $\nu_a = 1 + 4\pi\kappa_a$. In the case there $\kappa_a \ll 1$ we have $\chi_a \simeq \kappa_a$. Substituting these relations into the system of equations (2), (3), (4) and (7) and neglecting by nonlinear terms, we obtain a system of linear homogeneous equations in partial derivatives with constant coefficients. Passing to the following representation for small perturbations δf

$$\delta f = f(\omega, k) exp(-\imath\omega t + \imath kx)$$

yields a homogeneous system of algebraic equations.

In the second case when we consider equilibrium spin-current the linear set of QHD equations splits on four independent sets. One of them contains δM_x, δM_y, δJ_{xx}, δJ_{yx}, the second one contains n, δv_x, δM_z, J_{zx}, the third set includes δJ_{xy}, δJ_{yy}, and fourth includes δJ_{xz}, δJ_{yz}. Two last sets have the same solution, which is $\omega = 2\gamma B_0/\hbar$. The second set give us following dispersion equation

$$\omega^2 + k^2\left(\frac{k\omega M_0}{mn_0}\frac{4\pi J_0^{zx}}{\omega^2 - \frac{\hbar^2 k^4}{4m^2}} + \frac{4\pi\gamma^2 n_0}{m}\right) = 0. \qquad (9)$$

We can see that non-zero equilibrium spin current leads to existence of the additional term in the dispersion equation which increases degree of equation up to four from two in the absence of the equilibrium spin-current.

$$\omega^4 + \omega^2\left(\frac{4\pi n_0\gamma^2 k^2}{m} - \frac{\hbar^2 k^4}{4m^2}\right)$$
$$+ \frac{4\pi k^3 M_0}{mn_0} J_0^{zx}\omega - \frac{4\pi n_0\gamma^2}{m}\frac{\hbar^2 k^6}{4m^2} = 0. \qquad (10)$$

In the absence of equilibrium spin-current we have

$$\omega^2 = -4\pi\gamma^2 n_0 k^2/m.$$

Thus there is new solution which describes faster damping solution.

It seems more reasonable to chouse $\Omega = \sqrt{m/(\pi n_0)}\omega/(2\gamma k)$ as dimensionless frequency. In this case we get equation (10) in the form

$$\Omega^4 + (1-\alpha)\Omega^2 + \beta\Omega - \alpha = 0, \qquad (11)$$

where $\alpha = \hbar^2 k^2/(16\pi m n_0\gamma^2)$ is the parameter describing contribution of the quantum Bohm potential, $\beta = kM_0 J_0^{zx}/\gamma^2 n_0^2$ is the parameter describing contribution of the equilibrium spin-current. The both parameters α and β depend on module of wave vector k. Thus this equation allows to get $\Omega(k)$ dependence. Equation (11) gives two solutions. One of them is stable solution, which dispersion presented on Fig. (1). The second solution of equation (11) shows an instability, which exists due to the existence of the equilibrium spin-current J_0^{zx}. This solution is presented on Fig. (2).



FIG. 1. (Color online) The figure describes the dispersion dependence of the stable collective excitation described by equation (11).



FIG. 2. (Color online) The figure describes the second solution of equation (11. This solution is unstable.) The dispersion dependence and corresponding instability increment of the collective excitation are presented.

IV. CONCLUSION

To get influence of the spin-current on dynamics of magnetized dielectrics we have derived equation of the spin-current evolution as a part of the set of the QHD equations.

We have obtained that existence of an equilibrium spin current, which space component directed parallel to the direction of the wave propagation, leads to existence of two new wave solution. One of these solutions is stable, where the frequency of oscillation ω is approximately proportional to the square of the wave vector $\omega \sim k^2$. The second solution reveals instability, which is caused by the equilibrium spin current.

---

* andreevpa@physics.msu.ru
† lsk@phys.msu.ru

[1] S. Takahashi, S. Maekawa, Journal of Magnetism and Magnetic Materials **272**, 1423 (2004).
[2] I. Zutic, J. Fabian, S. Das Sarma, Rev. Mod. Phys. **76**, 323 (2004).
[3] Jairo Sinova and Igor Zutic, Nature Materials **11**, 368 (2012).
[4] Dmytro Pesin and Allan H. MacDonald, Nature Materials **11**, 409 (2012).
[5] Ivan J. Vera-Marun, Vishal Ranjan and Bart J. van Wees, Nature Physics **8**, 313 (2012).
[6] Ron Jansen, Nature Materials **11**, 400 (2012).
[7] Wei Han, K.M.McCreary, K.Pi , W.H.Wang, YanLi , H.Wen, J.R.Chen, R.K.Kawakami , Journal of Magnetism and Magnetic Materials **324**, 369 (2012).
[8] K. Uchida, H. Adachi, T. An, H. Nakayama, M. Toda, B. Hillebrands, S. Maekawa, and E. Saitoh, Journal Of Applied Physics **111**, 053903 (2012).
[9] F. Pezzoli, F. Bottegoni, D. Trivedi, F. Ciccacci, A. Giorgioni, P. Li, S. Cecchi, E. Grilli, Y. Song, M. Guzzi, H. Dery, and G. Isella, Phys. Rev. Lett. **108**, 156603 (2012).
[10] Z. Rashidian, F. Kheirandish, Int. J. Theor. Phys. **51**, 1989 (2012).
[11] Richard T. Hammond, Applied Physics Letters **100**, 121112 (2012).
[12] Jeongsu Lee, Rafal Oszwaldowski, Christian Gothgen, and Igor Zutic, Physical Review B **85**, 045314 (2012).
[13] Yuan Li, Mansoor B. A. Jalil, and Seng Ghee Tan, Journal Of Applied Physics **111**, 093717 (2012).
[14] Rong Zhang, LongBai, Chen-LongDuan, Physica B **407**, 2372 (2012).
[15] M. J. Ma, M. B. A. Jalil, and Z. B. Siu, Journal Of Applied Physics **111**, 07C326 (2012).
[16] N. T. Bagraev, O. N. Guimbitskaya, L. E. Klyachkin, A. A. Kudryavtsev, A. M. Malyarenko, V. V. Romanov, A. I. Ryskin, I. A. Shelykh, A. S. Shcheulin, Physica C **893**, 470 (2010).
[17] J. Flipse, F. L. Bakker, A. Slachter, F. K. Dejene and B. J. van Wees, Nature Nanotechnology **7**, 166 (2012).
[18] L. S. Kuz'menkov, S. G. Maksimov, and V. V. Fedoseev, Theor. Math. Fiz. **126** 136 (2001) [Theoretical and Mathematical Physics, **126** 110 (2001)].
[19] P. A. Andreev, L.S. Kuz'menkov, arXiv:1109.3335 (accepted for publication in Int. J. Mod. Phys. B).
[20] Z. An, F. Q. Liu, Y. Lin, C. Liu, Sci. Rep. **2**, 388 (2012).
[21] Farkhad G. Aliev, Juan F. Sierra, Ahmad A. Awad, Gleb N. Kakazei, Dong-Soo Han, Sang-Koog Kim, Vitali Metlushko, Bojan Ilic, and Konstantin Y. Guslienko, Phys. Rev. B **79**, 174433 (2009).
[22] A. A. Awad, K. Y. Guslienko, J. F. Sierra, G. N. Kakazei, V. Metlushko, and F. G. Aliev, Appl. Phys. Lett. **96**, 012503 (2010).
[23] G. N. Kakazei, P. E. Wigen, K. Yu. Guslienko, V. Novosad, A. N. Slavin, V. O. Golub, N. A. Lesnik, and Y. Otani, Appl. Phys. Lett. **85**, 443 (2004).
[24] Ki-Suk Lee, Konstantin Y. Guslienko, Jun-Young Lee, and Sang-Koog Kim, Phys. Rev. B **76**, 174410 (2007).
[25] K. Y. Guslienko, R. H. Heredero, O. Chubykalo-Fesenko , Phys. Rev. B **82**, 014402 (2010).
[26] P. A. Andreev, L.S. Kuz'menkov, Moscow University Physics Bulletin **62**, N.5, 271 (2007).
[27] L. S. Kuz'menkov and S. G. Maksimov, Teor. i Mat. Fiz., **118** 287 (1999) [Theoretical and Mathematical Physics **118** 227 (1999)].
[28] P. A. Andreev, L. S. Kuz'menkov, Phys. Rev. A **78**, 053624 (2008).
[29] M. I. Trukhanova, Int. J. Mod. Phys. B, **26**, 1250004 (2012).
[30] P. A. Andreev, L. S. Kuz'menkov, M. I. Trukhanova, Phys. Rev. B **84**, 245401 (2011).
[31] G. Brodin, M. Marklund, J. Zamanian, B. Ericsson and P. L. Mana, Phys. Rev. Lett. **101**, 245002 (2008).
[32] P. A. Andreev, L. S. Kuz'menkov, Physics of Atomic Nuclei **71**, N.10, 1724 (2008).
[33] P. A. Andreev and L. S. Kuz'menkov, PIERS Proceedings, p.1047, March 20-23, Marrakesh, MOROCCO 2011.
[34] L. S. Kuz'menkov, S. G. Maksimov, and V. V. Fedoseev, Vestn. Mosk. Univ., Ser. 3: Fiz., Astron., No. 5, 3 (2000) (Moscow Univ. Phys. Bull., No. 5, 1 (2000)].
[35] M. Marklund and G. Brodin, Phys. Rev. Lett. **98**, 025001 (2007).
[36] S. M. Mahajan and F. A. Asenjo, Phys. Rev. Lett. **107**, 195003 (2011).
[37] F. Haas, G. Manfredi, M. Feix, Phys. Rev. E **62**, 2763(2000).
[38] P. K. Shukla, B. Eliasson, Rev. Mod. Phys. **83**, 885 (2011).
[39] P. A. Andreev and L. S. Kuz'menkov, PIERS Proceedings, p. 1055, august 19-23, Moscow, Russia 2012.
[40] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. **67**, 1822 (1998).
[41] T. L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
[42] P. A. Andreev, arXiv: 1201.0779.
[43] P. A. Andreev and L. S. Kuz'menkov, arXiv:1208.1000.
[44] P. A. Andreev, arXiv: 1201.6553.